Conference Program
Day 3: Tuesday, May 25, 2021

12:00-12:10	**Welcome & Announcements** Edmund Walsh, PhD Student and Conference Planning Committee Member Arthur Labatt Family School of Nursing, Western University	
12:15-1:30	**Concurrent Session D: Oral Paper Presentations**	
	D1 – Violence Against Women: Considerations for Health & Social Services (Moderator: Logan Kennedy)	
	12:15-12:30	**Violence Against Women Services in a Pandemic: A Multi-Method Research & Knowledge Mobilization Project** Nadine Wathen
	12:35-12:50	**Impacts of COVID-19 Related Changes in Income on Women Experiencing Intimate Partner Violence** Cara A. Davidson, Tara Mantler, Andrew M. Johnson
	12:55-1:10	**Association Between Intimate Partner Violence and Functional Gastrointestinal Disease and Syndrome Among Adult Women: Systematic Review** Ohud Banjar, Marilyn Ford-Gilboe, Deanna Befus, Bayan Alilyyani
	1:15-1:30	**Lifetime Prevalence of Emotional/Psychological Abuse Among Qualified Female Healthcare Providers**
	Chronic Disease & Pain Management (Moderator: Maureen Leyser)	
	12:15-12:30	**Effectiveness of Home-Based Cardiac Rehabilitation and Its Importance During COVID-19** Hannah Pollock, Anna Garnett
	12:35-12:50	**Understanding the Social Determinants of Health from the Standpoint of Patients: An Institutional Ethnography of Mental Health, Addictions and Poverty in the Lives of People with Chronic Pain** Fiona Webster, Laura Connoy
	12:55-1:10	**Exploring Patient Engagement and the Use of Opioids in Managing Chronic Pain: A Scoping Review** Bayan Alilyyani, Ryan Chan, Laura Connoy, Fiona Webster
	D3 – Promoting Practice & Education Across Diverse Contexts (Moderator: Cara Davidson)	
	12:15	

E3 – Science Pitch Session I (Moderator: Fiona Webster)

1:45-2:40

Harm Reduction Services in Ottawa: The Culture of Drug Use
Marlene Haines, Patrick O'Byrne

Opioid Crisis: A Qualitative Analysis of Financial Influences and Addiction
Nicole Naccarato, Noah Wacker, Lissa Gagnon

Mental Health and Addictions Care Provided by Nurses in the Emergency Department

3:00-4:00	**Concurrent Session F: Oral Paper Presentations**	
	F1 – Mental Health: Suicide & Schizophrenia (Moderator: Heather Sweet)	
	3:00-3:15	**Trends and Factors Associated with Suicide Deaths in Older Adults** Eada Novilla-Surette, Salimah Shariff, Britney Le, Richard Booth
	3:20-3:35	**Zero Suicide: St. Joseph's Health Care London and Beyond** Amy Van Berkum, Shauna Graf
	3:40-3:55	**Engagement, Partnership & Participation in Self-Management in Outpatient Services for People with Schizophrenia** Mary-Lou Martin, Susan Strong, Heather McNeely, Lori Letts, Alycia Gillespie
	F2 – Violence in the Workplace (Moderator: Edmund Walsh)	
	3:00-3:15	**Putting the Brakes on Aggressive Behaviours: Empowering Nurses Using the "Traffic Light Process"** Chantal Singh, Karen Laidlaw

ABSTRACTS

Concurrent Session D: Oral Paper Presentations

Impacts of COVID-19 Related Changes in Income on Women Experiencing Intimate Partner Violence

Cara A. Davidson, School of Health Studies, Western University, London, Ontario
Tara Mantler, School of Health Studies, Western University, London, Ontario
Andrew M. Johnson, School of Health Studies, Western University, London, Ontario
a-1 (rnm)-41 (tm4 (o6 (t)Tj qC uC aEM- (a4 (,)1-1 (rnf(it)-- (e)13 (w (e)9(L)41.66 (ha)7 (y (t)-)th O)3 (n)-(h)1w 0.31(e)1

Lifetime Prevalence of

D2 – Chronic Disease & Pain Management

Effectiveness of Home-Based Cardiac Rehabilitation and Its Importance during COVID-19

Hannah Pollock, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Anna Garnett, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Background: Cardiac rehabilitation is a critical disease management and health promotion intervention for individuals recovering from cardiac events and living with heart disease, proven to reduce the risk of mortality, morbidity, and hospitalizations. The current COVID-19 pandemic has caused disturbances in every aspect of life including the postponement or cancellation of many health services, including 41% of cardiac rehabilitation programs in Canada. Without access to cardiac rehabilitation, individuals face many challenges trying to recover from cardiac events and improve their health and well-being. Education, resources and support normally provided in a traditional centre-based program are limited due to COVID-19 closures, placing participants in a potentially risky situation to suffer adverse health outcomes.

Purpose: This position suggests that transitioning from a centre-based cardiac rehabilitation program to a home-based program during the COVID-19 pandemic and the foreseeable future is a more effective health strategy to provide continuous care to cardiac patients.

Evidence: Home-based cardiac rehabilitation programs have been utilized in Canada prior to the pandemic but vary in structure and function. They are typically defined as those in which the majority of exercise is performed without direct, hands-on, line-of-sight supervision. Programs utilize a variety of resources, including technology, to regularly monitor participants and allow them to engage in education and counselling resources remotely. The program's flexibility and convenience overcome many of the multi-level barriers which normally impede participants from accessing centre-based programs. Home-based programs have proven to be equally effective, if not more effective than centre-based programs in improving mortality, cardiac events, exercise capacity and modifiable risk factors.

Implications: Home-based cardiac rehabilitation programs are a valid alternative to support and protect a vulnerable population, especially those at high risk of complications if diagnosed with COVID-19. Transitioning to a home-based platform may be a challenge, however, home-based rehabilitation is better

Understanding the Social Determinants of Health from the Standpoint of Patients: An Institutional Ethnography of Mental Health, Addictions and Poverty in the Lives of People with Chronic Pain

Fiona Webster, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Laura Connoy, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Chronic pain (CP) is a significant health problem and is the most prevalent chronic condition in Canada. Increased rates of opioid prescribing have followed this high prevalence of CP, contributing to an increase in opioid-related deaths. Based on our previous research, which examined CP care in Ontario from the perspective of primary care providers, many patients who suffer from CP also suffer from poverty, poor mental health, and addiction –

Exploring Patient Engagement and the Use of Opioids in Managing Chronic Pain: A Scoping Review

Bayan Alilyyani, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Enhancing Advanced Practice Nursing: The Value of Role Clarity and Mentorship

Lisa Morgan, St. Joseph's Health Care London, London, Ontario
Alexis Smith, St. Joseph's Health Care London, London, Ontario
Amanda Thibeault, St. Joseph's Health Care London, London, Ontario

Problem Statement: Advanced practice nurses (APN), defined by clinical nurse specialist and nurse practitioner roles, have become vital to meet the needs of our growing health care system (Canadian Nurses Association, 2019). There are several documented challenges faced when implementing APN roles, including lack of role clarity and professional boundary-setting, unsupportive leadership, and reduced job satisfaction, which contribute to variable role optimization and retention (Contandriopoulos et al. 2015; Kilpatrick et al. 2016). Evidence supports that successful APN role implementation requires strong support from operational leadership, and formalized on-boarding processes including a comprehens (d)-4 (o)8 di(d)6-41 (ad)-12.1 2 (n)-4 (t)-50

Association, 2eng anototzat (sH)o ve Auictace-2 (me) (lar-3.9 (io)-2 (n)-4 (,)11 6(2)-2 (C

The Challenge and Potential of Trauma- and Violence-informed Care for Nurses Working with Women who have Experienced Intimate Partner Violence

Noël Patten Lu, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Marilyn Ford-Gilboe, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Lorie Donelle, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Victoria Smye, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Kimberley Jackson, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Background and Purpose: Intimate partner violence (IPV), one of the most common forms of violence against women, is a serious public health issue that affects 1 in 4 women in Canada. As the first and possibly only point of contact in the healthcare system for women who have experienced IPV, nurses can have profound effects (positive and negative) on women's health, safety, and how they engage with health services. However, healthcare providers, including nurses, often lack appropriate training to deal with IPV, which can contribute to further harms and re-traumatization. [illegible]

D4 – Mental Health & Homelessness

From Hospital to Homelessness: Preventing Discharge to "No Fixed Address"v2

Cheryl Forchuk, Lawson Health Research Institute; Arthur Labatt Family School of Nursing, Western University, London, Ontario
Jeffrey Reiss, London Health Sciences Center, London, Ontario
Sherri Lawson, London Health Sciences Centre, London, Ontario
Sandra Northcott, St. Joseph's Health Care London, London, Ontario
Rebecca Vann, St. Joseph's Health Care London, London, Ontario
Dan Catunto, Ontario Works Operations, City of London, London, Ontario
Sarah Collins, Salvation Army Centre of Hope, London, Ontario
Richard Booth, Arthur Labatt Family School of Nursing, Western University, London, Ontario

The discharge of psychiatric patients from the hospital into homelessness is a prevalent issue despite research indicating social, safety, health and economic detriments on both the individual and community level (Forchuk et al., 2008; Gaetz, 2012). Lack of stable housing for discharged inpatients results in long-term consequences including exacerbation of health problems and costly health care service use and hospital readmission (Mikkonen & Raphael, 2010; Munn-Rivard, 2014). Patients experiencing homelessness are four times more likely to be readmitted within a month of discharge and hospitals must spend $2,559 more per client (Hwang et al., 2011; Saab et al., 2016). Finding safe housing for these individuals is imperative to their recovery and transition back to the community.

The "No Fixed Address" version 2 (NFA v.2) project has tested the efficacy of a potential best practice program that finds safe housing for inpatients, preventing discharge from hospital into homelessness. Forchuk and colleagues developed a system that streamlines housing and social supports using on-site access. Housing Stability Workers and Ontario Works are brought directly into hospital, allowing inpatient access via drop-in or by appointment.

Findings of the NFA v.2 project will be discussed. In a previous pilot project, Forchuk et al., (2006) found that all seven participants randomly assigned to the intervention remained housed at 3 and 6 months' follow-up, while individuals in usual care were unhoused or had entered the sex trade. In a following scaled-up phase of the project involving 219 acute psychiatric clients and 32 tertiary care clients (Forchuk et. al, 2008), 92.5% of those who accessed the service and were at risk of homelessness were connected with affordable accommodation.

Since homelessness has a detrimental effect on recovery, client and community safety, and healthcare expenditure, locating safe housing for psychiatric patients may have a positive impact on treatment, rehabilitation, and the system as a whole. The findings of this project may offer safe policy alternatives for the prevention of homelessness for at risk individuals.

Mental Health Experiences and Pathways to Homelessness Among Refugee Claimants

Bridget Annor

Factors Influencing Access and Utilization of Health Service among Substance Using Homeless Youth: A Scoping Review

Vanisa Ezukuse, Arthur Labatt Family School of Nursing, Western University, London, Ontario

In Canada an estimated 35,000 to 40,000 youth (ages 13-24) experience homelessness yearly (Gaetz et al., 2014). As a diverse population with numerous stressors, stigmatization transient housing, homeless youth often use substances as a form of coping. Despite the higher rates of physical illness and mental disorders prevalent among homeless youth they have been shown to have lower utilization of and access to healthcare. (Abel-baki et al., 2019; Karabanow et al., 2018). As a result, it is critical to recognize and understand the factors that influence service use among this population.

The purpose of this paper is to: (1) Review the literature on substance using homeless youth's access to health care services, (2) Identify and synthesize factors that influenc

A Scoping Review of Promising Structural Reforms to Support Youth Mental Health

Abe Oudshoorn, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Michelle Virdee, Western University, London, Ontario
Joseph Adu, Western University, London, Ontario
Ross Norman, Western University, London, Ontario
Eugenia Canas, Western University, London, Ontario
Romaisa Pervez, Western University, London, Ontario
Arlene MacDougall, Parkwood Institute, London, Ontario

The purpose of this presentation is to share the findings of a scoping review on structural reforms to support youth mental health. The mental health experiences and needs of transitional age youth has been identified as a priority both for research and for clinical interventions. While promising programs are being enacted globally, these are often delivered in the context of systems that are not functioning optimally, thus being a form of down-stream response. Therefore, true transformation of youth mental health supports needs to come for structural reforms rather than simply "working around the margins". As part of a larger social innovation lab aimed at shifting mental health outcomes for transitional age youth, this scoping review was designed to provide an evidence-based foundation for considering structural reforms. The search included EMBASE, CINAHL, PsychInfo, Medline, Socialogical Abstracts, Proquest Dissertations and Theses, and the Nursing and Allied Health Database. Search terms focused on interventions at the structural, system, or policy level, anywhere in the world, and including both empirical and theoretical work. A team of 6 researchers worked through an initial pool of 5,652 articles through title, abstract, and full text screening. Ultimately, 61 articles were included in the data extraction with priority put on articles that scored high in a rating of congruence, relevance, innovation, feasibility, and clarity. This presentation will focus on key findings from the review, highlight potential opportunities for structural reforms in a Canadian context to enhance youth mental health.

Concurrent Session E: Oral Paper Presentations & Science Pitch Sessions

E1 – Digital Health: Smartphones & Mobile Applications

Evolving a Personalized, Online Safety & Health Resource for Women Experiencing Intimate Partner Violence to a Publicly Available App: My Plan Canada

Using Snapchat to Promote STI Screening at a Rural Public Health Unit

Brad Hiebert, Huron County Health Unit
Annie O'Dette, Huron County Health Unit
Marian Doucette, Huron County Health Unit
Rita Marshall, Huron County Health Unit
Chisomo Mchaina, Huron County Health Unit
Kate Underwood, Huron County Health Unit

BACKGROUND & OBJECTIVES: Since 2013 in Huron County (a predominantly rural region), males under 30

Nurses' Use of Smartphones and Mobile Phones in the Workplace: A Scoping Review

Andrea de Jong, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Lorie Donelle, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Michael S. Kerr, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Introduction: There has been a significant increase in technological infrastructure of many healthcare organizations to support the practice of healthcare providers; however, many nurses are using their personal digital devices, smartphones and mobile phones while at work for personal and professional purposes. Despite the proliferation of smartphone use in the healthcare setting, there is limited research available on the clinical use of these devices by nurses. This study aimed to understand the current breadth of research on nurse's personal smartphone use in the workplace and identify implications for research, practice, and education.

Methods: A scoping review was conducted and the following databases were used in the literature search: CINHAL, PubMed, Dissertations and Theses, EMBASE, MEDLINE, Nursing and Allied Health Database, Scopus, Web of Science, and Cochrane Reviews. Search terms used were: Nurs* AND (personal digital technology OR smartphone OR cellphone OR mobile phone OR cellular phone).

Results: Sixteen of 1765 articles met inclusion criteria. All but three articles focused on personal device use in acute care settings. Three main themes from the thematic analyses of the reviewed literature included: personal smartphone use for patient care, personal smartphone use for personal reasons, and implications of personal smartphone use. Nurses used their smartphones to locate information about medications, procedures, diagnoses, and laboratory tests. Nurses reported improved communication between health team members and used their personal devices to communicate patient information via text messaging, calling, and picture/video functions. Yet, nurses expressed insight into personal smartphone use and challenges related to distraction, information privacy, organizational policies, and patient perception.

Conclusion: Nurses are bringing their smartphones to work and see these devices as an efficient method to gather patient care information and to communicate with the healthcare team. This review highlights knowledge gaps regarding nurses' personal device use and information safety, patient care outcomes, and communication practices among healthcare teams. The current breadth of research is focused on acute care, with little to no research focus in other practices settings. Research initiatives are needed to explore personal device use across the continuum of workplace settings, such as homecare, long-term care, and public health.

The Environment as Patient: A Content Analysis of Canadian Nursing Organizations and Regulatory Bodies Policies on Nurses' Role in Environmental Health

Courtney Allen, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Lorie Donelle, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Individual, community, and societal health is impacted by the environment which is impacted by pollution of air, water and soil, and climate change. Poor environmental health conditions have been associated with

E3 – Science Pitch Session I

Harm Reduction Services in Ottawa: The Culture of Drug Use

Marlene Haines, School of Nursing, University of Ottawa, Ottawa, Ontario
Patrick O'Byrne, School of Nursing, University of Ottawa, Ottawa, Ontario

Mental Health and Addictions Care Provided by Nurses in the Emergency Department

Shubhjit Gabhi, Arthur Labatt Family School of Nursing, Western University, London, Ontario, Canada

Seminal pieces of literature from various countries throughout the world including Canada, Australia and the United States of America report that Emergency Departments (ED) are not suitable for people with specialized Mental health and addictions (MHA) needs and that ED nurses do not feel well-equipped to care for this patient population (Fleury et al., 2020; Innes et al., 2014; Marynowski-Traczyk & Broadbent, 2011; Reed & Fitzgerald, 2005; Thomas et al., 2018; Wolf et al., 2015; Zolnierek & Clingerman, 2012; Zun, 2012). A valid reason as to why nurses feel this way may be related to a gap in nursing education and training. Many nursing regulators in Canada including the College of Nurses of Ontario take a generalist approach to nursing education required to practice as a Registered Nurse (RN) or Registered Practical Nurse (RPN) (College of Nurses of Ontario, 2019). Generalist prepared nurses have the knowledge, skill and judgement to provide safe care to people of all ages and genders in a wide variety of practice settings (CNO, 2019). This means that nursing students receive some exposure to theoretical knowledge and clinical practice in the area of MHA; yet this may not be sufficient in preparing future nurses to care for this population in the settings where they are now receiving care. I propose enacting a qualitative research methodology to engage emergency nurses to describe their mental health practices. Recognizing that nursing curricula are already packed, the goal of this research will be to develop practical, structured education and training programs for ED nurses that can be completed by practicing nurses.

Investigating Physiological Determinants of Mental Health in Children with Cerebral Palsy

Daniela Testani, Western University, London, Ontario
Laura Brunton, Western University, London, Ontario
Carly McMorris, University of Calgary, Calgary, Alberta

Background
Fifty-seven percent of children with cerebral palsy (CP) experience mental health symptoms including anxiety and depression.1 Although CP is non progressive, the development of secondary conditions can have progressive effects on an individuals' functional abilities. Particularly, untreated mental health symptoms can have a negative effect on children's quality of life. Children with CP are more likely to experience fatigue, chronic pain and sleep disturbances. Identifying modifiable risk factors that contribute to the progression of depressive and anxiety symptoms can be vital in preventing lifelong challenges into adulthood. While it is theorized that the combined presence of these secondary symptoms may worsen the effects of mental health, that is yet to be systematically examined.

Objectives and Hypotheses
My objective is to better understand the relationship between fatigue, pain and sleep on mental health symptoms for children with CP. In addition, as a secondary objective we are observing the relationship between physical activity and its association with mental health. We hypothesize that moderate to severe levels of pain and/or fatigue, as well as sleep difficulties will be associen (s)2 (le)-1(ie)-1 (s)nt6 (e)1 (snn)10 (t6 5e)1 fn

Implementing Breastfeeding Education in Pediatric Settings

Keri Durocher, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Jody Ralph, School of Nursing, University of Windsor, Windsor, Ontario

Breastfeeding adherence rates in community and acute hospital settings are substandard across many developed nations, despite the development of programs to support them. For example, the Baby-friendly Hospital Initiative program was developed by the World Health Organization to enhance breastfeeding success. Based on a narrative review of the literature relevant to community and acute pediatric healthcare settings, it is evident that enhanced education for interdisciplinary team members needs to be implemented to support breastfeeding. Findings from twenty-eight articles are included in this practice initiative, including systematic reviews, randomized control trials, case control, cohort, descriptive, and qualitative studies, as well as opinion articles. After synthesizing the study results into content themes, it is evident that initiation and management of breastfeeding within these settings can be improved through increased healthcare provider knowledge. A narrative summary of the evidence reveals that issues related to breastfeeding promotion in community and acute pediatric settings are due to complacency with early cessation, inadequate healthcare provider knowledge, and overreliance on Internationally Board-Certified Lactation Consultants® (IBCLCs). Innovative recommendations from gathered studies include educational approaches that can be implemented through managerial and clinical strategies, structured breastfeeding education programs, and module-based learning. Hands-on learning of healthcare professionals with assistive devices to troubleshoot breastfeeding issues will also assist with breastfeeding success in pediatric settings. If pediatric healthcare leaders can adopt the outlined strategies to their organizational needs, breastfeeding success will be enhanced in the future.

Meditating in Virtual-Reality: Investigating Affect Responses of Mindfulness through a Trauma-Informed and Instructor Present Approach

Madison Waller, Department of Psychology, Western University, London, Ontario
Paul Frewen, Departments of Psychiatry and Psychology, Western University, London, Ontario

Mindfulness meditation (MM) as a therapeutic intervention is increasingly being investigated as a possible treatment for a variety of psychological disorders, including depression, anxiety, and most prominently, posttraumatic stress disorder (PSTD). One fo

Intersectoral Collaboration: A Literature Review

Patrick Ellis, Arthur Labatt Family School of Nursing, Western University, London, Ontario

With the introduction of the Ontario Health Teams model, there exists a need to understand the level of importance that teamwork will have across healthcare agencies. The new model is proposed to introduce improved continuity of care with increased patient involvement. Although there is discussion over implementation of this system, organizations have not focused on how teamwork surrounding intersectoral collaboration -8 (e)1T((ns)6 (h)1llab)-4 (o)-2 (ra)18 (e)d(h)-4 (u)rah-4 (i1.1 (t)-4 (.)o)-2 t-4 (io)-2 (n) (o)-2 oration -8 (

Examining the Impact of Managers' Authentic Leadership on Long-Term Care Nurses' Job Turnover Intentions

Edmund J. Walsh, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Michael S. Kerr, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Carol A. Wong, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Emily A. Read, Faculty of Nursing, University of New Brunswick, Fredericton, New Brunswick
Joan Finegan, Faculty of Social Science, Western University, London, Ontario

Purpose: The purpose of this p

The Context and Consequences of Being Turned Away from a Domestic Violence Shelter

Rachel Colquhoun, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Domestic violence (DV) shelters provide a wide range of supports for women experiencing violence, yet fewer than 20% of Canadian women access these services. Many do seek help from shelters but are turned away. Women's Shelters Canada (2019) reports that the turn-away rate in Canada is 79% and current research does not capture the unique challenges of women who are turned away. Little is known about the characteristics of women who are turned away; the context that shapes this experience; or the consequence for their health, well-being and safety. The purpose of this study is to understand variations in women's experiences of being turned away from a DV shelter in the context of shelter seom a (sTw -8 (n)-4 8(e)3 (l)i ()4 (o)1 (te)3 (r)4 (y)3

Health-Seeking Behaviour Related to Selected Dimensions of Wellness in Community Dwelling Older Adults

Navjot Gill, Health and Rehabilitation Sciences Program, Western University, London, Ontario
Denise Connelly, School of Physical Therapy, Western University, London, Ontario

Background
Older people generally prefer to ‘stay-put' in their own home. Informed by the Seven Dimensions of Wellness, a component of the International Council on Active Ageing (ICAA) Model, measures of physical function (i.e. physical), fall risk (i.e. environmental), and psychosocial factors (i.e. emotional, spiritual and social) related to health have been selected with the assumption that they intersect to influence health-seeking behaviour of older adults and ageing-at-home.

Objective and Hypothesis
The objective of the study is to investigate the relationship among dimensions of well-being, including physical function, fall risk, psychosocial factors and awareness of community-support services, with health-seeking behaviour in community-dwelling older adults.

Null Hypothesis
The scores obtained on health dimension outcomes will not predict health-seeking behaviour in community-dwelling older adults

Alternate Hypothesis
The scores obtained on health dimension outcomes will predict health-seeking behaviour in community-dwelling older adults.

Proposed Methods
This project will use correlational design. A sample size of 98 subjects will be recruited. Subjects will be older

(c)86 (o)fu
in9 (o-4 (l me)-1 u9 (olt)6 (e)-1 (ig) (o)-2 (9)-2-1 (rlin9 (o 28.58 (S)101 (d)-4 g (i)14 0.9 (s)2 (amp)-2 (i)]TJ)10 (e)-1 (

A Narrative Review of Post-Trauma Resilience and Optimism Frameworks, and Proposal of an Integrated Framework for Musculoskeletal Trauma

Wonjin Seo, Faculty of Health Sciences,

Zero Suicide: St. Joseph's Health Care London and Beyond

Amy Van Berkum, St. Joseph's Health Care London, London, Ontario
Shauna Graf, St. Joseph's Health Care London, London, Ontario

St. Joseph’s Health Care London (St. Joseph’s) is leading the way in the implementation of Zero Suicide across Canada. On average, ten Canadians die by suicide each day, making suicide the ninth leading cause of death
i

Engagement, Partnership & Participation in Self-Management in Outpatient Services for People with Schizophrenia

Mary-Lou Martin, St. Joseph's Healthcare Hamilton/McMaster University, Hamilton, Ontario
Susan Strong, St. Joseph's Healthcare Hamilton/McMaster University, Hamilton, Ontario
Heather McNeely, St. Joseph's Healthcare Hamilton/McMaster University, Hamilton, Ontario
Lori Letts, McMaster University, Hamilton, Ontario
Alycia Gillespie, St. Joseph's Healthcare Hamilton, Hamilton, Ontario

Project Goal & Description: To operationalize, implement and evaluate SET for Health, a model of self-management support within the case management care process of specialized outpatient services for people with schizophrenia.

Methods: A mixed method feasibility study examined to what extent SET for Health: added value from clients' and providers' perspectives; and influenced individuals' engagement in treatment, participation in self-management, symptom distress, hope, and quality of life. Facilitation tools/procedures operationalize the intervention derived from standardized self-management programs. 10 case managers (registered nurses, social workers, occupational therapists) are delivering the intervention to 50 clients.

Phases of the project: Clients, families and staff are involved in the evaluation and giving recommendations.

Results: At 75% retention, SET for Health offered mechanisms for client participation, engagement and voice; self-management discussions of illness and health; and provider recovery orientation. Delivery variations were noted across case managers related to challenges in changing usual practices, particularly client directed goal-setting, problem-solving and review.

Lessons Learned: Self-management support can be delivered and benefit clients with complex living challenges. Support (institutional, education, environmental) was needed to integrate self-management into routine care. The ability to engage and work together with clients and deliver an accessible, feasible model of self-management support valued by clients and providers, coupled with Health Quality Ontario's 2018 standards and Lean and colleagues meta-analysis (2019) findings make a case for this type of partnership and routine delivery. Self-management commands attention as an intervention option for high risk clients with limited insight, negative symptoms and medical co-morbidities. Significant pre-post paired measures of self-management, social and occupational functioning, and illness severity need replication in a controlled study.

Sponsors: Dr. Ian & Shirley Rowe Research Award, Research Institute of St. Joseph`s Healthcare Hamilton.

Part of the Job? Gender as a Determinant of Workplace Violence Against Nurses

Andrea Baumann, University of Toronto, Lawrence S. Bloomberg Faculty of Nursing, Toronto, Ontario
Sioban Nelson, University of Toronto, Lawrence S. Bloomberg Faculty of Nursing, Toronto, Ontario

The problem of workplace violence against nurses is well documented in the academic and policy literatures. However, existing studies tend to conceptualize violence as a problem between individual health care workers and patients, largely failing to examine underlying power structures that may be contributing to

Implementing a Workplace Violence Reporting System for Nurses in a Healthcare Setting in Pakistan

Rozina Somani, Lawrence S. Bloomberg, Faculty of Nursing, University of Toronto, Toronto, Ontario
Carles Muntaner, Lawrence S. Bloomberg, Faculty of Nursing and Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario
Edith Hillan, Lawrence S. Bloomberg, Faculty of Nursing, University of Toronto

F3 – Understanding Health: A Global Perspective

Nurses and Midwives' Experience of Providing Fertility Awareness-based Methods including Natural Family

The Correlation be

A Feminist Narrative Inquiry into Being a Child Bride in Nigeria

E-health Decision Support Technologies in the Prevention and Management of Pressure Ulcers: A Systematic Review

Justine Ting, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Anna Garnett, Arthur Labatt Family School of Nursing, Western University, London, Ontario

Background: Pressure ulcers are problematic across clinical settings, negatively impacting patient outcomes while resulting in substantial costs to the healthcare system. E-health clinical decision support technologies can play a key role in improving pressure ulcer-related outcomes.

Purpose: The aim of this systematic review was to assess the impact of e-health decision support interventions on pressure ulcer management and prevention.

Methods: This review utilized the systematic review protocol outlined by the Joanna Briggs Institute Manual for Evidence Synthesis. The research question guiding this inquiry was: What is the impact of e-health decision support technologies on the prevention and management of pressure ulcers? This systematic review located 19 studies from January 2010 to October 2020 and identified a range of e-health clinical decision support technologies.

Results: Most interventions were integrated in electronic health records and were implemented primarily in long-term care and hospital settings. E-health clinical decision support interventions performed a variety of functions, including: generating tailored recommendations for care planning, creating automated pressure ulcer reports, and providing cues to promote adherence to practice standards. Image-processing software to generate measurement and staging recommendations was also identified as an emerging area of research. The findings of this review revealed promising results regarding the usability and accuracy of e-health clinical decision support tools. Results of the review also indicated improved adherence to pressure ulcer prevention practices and clinician staging accuracy. However, the studies included in this review did not consistently show reductions in pressure ulcer prevalence, incidence, or risk.

Implications: More high-quality studies are needed to establish the types of e-health clinical decision support tools that can drive sustainable improvements to pressure ulcer-related patient outcomes. These may include randomized controlled trials, large-scale studies spanning multiple institutions, or targeted studies exploring the applications of e-

“You have to be careful”: Examining Children’s Perspectives Related to Digital Device and Social Media Use through a Digital Health Lens

Danica Facca, Faculty of Information and Media Studies, Western University, London, Ontario
Lorie Donelle, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Shauna Burke, School of Health Studies, Western University, London, Ontario
Bradley Hiebert, Arthur Labatt Family School of Nursing, Western University, London, Ontario
Emma Bender, School of Kinesiology, Western University, London, Ontario
Stephen Ling, Faculty of Science, Western University, London, Ontario

Children are becoming regular users of digital devices and, by extension, social media. This pilot study used a cross sectional survey design to explore how 42 young children (aged six to ten years) in Ontario, Canada perceived their access to, use of, and privacy associated with digital device and social media at home and school. From a digital health perspective, it is important for parents, educators, and researchers to understand children’s digital practices in order to best support their learning, growth, and wellbeing within the digital age.